arXiv:0808.0377v1 [math.GR] 4 Aug 2008

CHARACTERIZATION OF $\mathrm{SL}(2,q)$ BY ITS NON-COMMUTING GRAPH

ALIREZA ABDOLLAHI

ABSTRACT. Let G be a non-abelian group and $Z(G)$ be its center. The non-commuting graph $\mathcal{A}_G$ of G is the graph whose vertex set is $G\backslash Z(G)$ and two vertices are joined by an edge if they do not commute. Let $\mathrm{SL}(2,q)$ be the special linear group of degree 2 over the finite field of order q. In this paper we prove that if G is a group such that $\mathcal{A}_G \cong \mathcal{A}_{\mathrm{SL}(2,q)}$ for some prime power $q \geq 2$, then $G \cong \mathrm{SL}(2,q)$.

1. Introduction and Results

Let G be a non-abelian group and $Z(G)$ be its center. One can associate with G a graph whose vertex set is $G\backslash Z(G)$ and two vertices are joined by an edge whenever they do not commute. We call this graph the non-commuting graph of G and it will be denoted by $\mathcal{A}_G$. The non-commuting graph $\mathcal{A}_G$ was first introduced by Paul Erdös [4] to formulate the following question: If every complete subgraph of $\mathcal{A}_G$ is finite, is there a finite bound on the cardinalities of complete subgraphs of $\mathcal{A}_G$? Neumann [4] answered positively Erdös question by proving that $|G : Z(G)| = n$ is finite and n is obviously the requested finite bound.

The non-commuting graph has been studied by many people (see e.g., [1],[3] and [5]). It is proved in [7] (resp., in [8]) that if G is a finite group with $\mathcal{A}_G \cong \mathcal{A}_{\mathrm{PSL}(2,q)}$ (resp., $\mathcal{A}_G \cong \mathcal{A}_{A_{10}}$), then $G \cong \mathrm{PSL}(2,q)$ (resp., $G \cong A_{10}$). For any prime power q, let $\mathrm{GL}(2,q)$ (resp. $\mathrm{SL}(2,q)$) be the general (resp. special) linear group of degree 2 over the finite field of order q. In this paper we study the groups whose non-commuting graphs are isomorphic to either $\mathrm{GL}(2,q)$ or $\mathrm{SL}(2,q)$. Our main results are the following.

Theorem 1.1. *Let G be a group such that $\mathcal{A}_G \cong \mathcal{A}_{\mathrm{GL}(2,q)}$ for some prime power $q > 3$. Then $G/Z(G) \cong \mathrm{PGL}(2,q)$, $G' \cong \mathrm{SL}(2,q)$ and $Z(G)$ is of order $q-1$. In particular, if q is even, then $G = G' \times Z(G)$.*

Theorem 1.2. *Let G be a group such that $\mathcal{A}_G \cong \mathcal{A}_{\mathrm{SL}(2,q)}$ for some prime power $q \geq 2$. Then $G \cong \mathrm{SL}(2,q)$.*

For any prime power q, we denote by $\mathrm{PGL}(2,q)$ (resp. $\mathrm{PSL}(2,q)$) the projective general (resp. special) linear group of degree 2 over the finite field of order q.

1991 *Mathematics Subject Classification.* 20D60.

Key words and phrases. Non-commuting graph; General linear group; Special linear group.

Author's Address: Department of Mathematics, University of Isfahan, Isfahan 81746-73441, Iran; and School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O.Box: 19395-5746, Tehran, Iran. e-mail: `abdollahi@member.ams.org`.

This research was in part supported by a grant from IPM (No. 87200118).

2. **Proofs**

Here for convenience, we remind some of the properties of non-commuting graphs and common properties of groups with isomorphic non-commuting graphs.

Let G and H be two non-abelian groups such that $\mathcal{A}_G \cong \mathcal{A}_H$. By Lemma 3.1 of [1], if one of G or H is finite, then so is the other. The order of $\mathcal{A}_G$ is $|G| - |Z(G)|$ and so $|G| - |Z(G)| = |H| - |Z(H)|$. The degree of a vertex x in $\mathcal{A}_G$ is equal to $|G| - |C_G(x)|$. Thus the multisets of degrees of vertices of two graphs $\mathcal{A}_G$ and $\mathcal{A}_H$ are the same.

A non-abelian group G is called an AC-group, if the centralizer $C_G(x)$ of every non-central element x of G is abelian.

Recall that a non-empty subset X of the vertices of a simple graph Γ is called independent if every two distinct vertices of X are not joint by an edge in Γ. Thus an independent set S of the non-commuting graph of a group is a set of pairwise commuting non-central elements of the group.

Lemma 2.1. *Let G and H be two finite non-abelian group with $\mathcal{A}_G \cong \mathcal{A}_H$.*

(1) *If $|G| = |H|$, then the multisets (sets with multiplicities) $\{|C_G(g)| : g \in G\backslash Z(G)\}$ and $\{|C_H(h)| : h \in H\backslash Z(H)\}$ are equal.*

(2) *If G is an AC-group, then H is also an AC-group.*

Proof. (1) It is straightforward, if we note that the set of non-adjacent vertices to a vertex x in the non-commuting graph H is $C_H(x)\backslash Z(H)$, and note that from $|G| = |H|$ we also have $|Z(G)| = |Z(H)|$, since $|H| - |Z(H)| = |G| - |Z(G)|$.

(2) Note that a subgroup S of a non-abelian group K is abelian if and only if either $S\backslash Z(S)$ is empty or $S\backslash Z(S)$ is an independent set in the non-commuting graph $\mathcal{A}_K$. Let ϕ be a graph isomorphism from $\mathcal{A}_H$ onto $\mathcal{A}_G$. Then it is easy to see that for each $h \in H\backslash Z(H)$,

$$C_H(h)\backslash Z(H) = \phi^{-1}\big(C_G(\phi(h))\backslash Z(G)\big). \qquad (*)$$

Now since G is an AC-group, $C_G(g)$ is abelian for all $g \in G\backslash Z(G)$ and so it follows from $(*)$ and the remark above that $C_H(h)$ is abelian. Hence H is also an AC-group. $\square$

Finite non-nilpotent AC-groups were completely characterized by Schmidt [6]. We use the following results in our proofs.

Theorem 2.2. ([6, Satz 5.9.]) *Let G be a finite non-solvable group. Then G is an AC-group if and only if G satisfies one of the following conditions:*

(1) *$G/Z(G) \cong \mathrm{PSL}(2,p^n)$ and $G' \cong \mathrm{SL}(2,p^n)$, where p is a prime and $p^n > 3$.*
(2) *$G/Z(G) \cong \mathrm{PGL}(2,p^n)$ and $G' \cong \mathrm{SL}(2,p^n)$, where p is a prime and $p^n > 3$.*
(3) *$G/Z(G) \cong \mathrm{PSL}(2,9)$ and G' is a covering group of A_6. In particular, G' is isomorphic to*

$$\begin{aligned}\mathcal{A} \cong < c_1, c_2, c_3, c_4, k \mid c_1^3 = c_2^2 = c_3^2 = c_4^2 = (c_1c_2)^3 = (c_1c_3)^2 = \\ = (c_2c_3)^3 = (c_3c_4)^3 = k^3, (c_1c_4)^2 = k, \\ c_2c_4 = k^3c_4c_2, kc_i = c_ik (i = 1, \ldots, 4), k^6 = 1 > .\end{aligned}$$

(4) *$G/Z(G) \cong \mathrm{PGL}(2,9)$ and $G' \cong \mathcal{A}$.*

For a finite simple graph Γ, we denote by $\omega(\Gamma)$ the maximum size of a complete subgraph of Γ. So $\omega(\mathcal{A}_G)$ is the maximum number of pairwise non-commuting elements in a finite non-abelian group G.

Theorem 2.3. (Satz 5.12 of [6]) *Let G be a finite non-abelian solvable group. Then G is an AC-group if and only if G satisfies one of the following properties:*

1. *G is non-nilpotent and it has an abelian normal subgroup N of prime index and $\omega(\mathcal{A}_G) = |N : Z(G)| + 1$.*
2. *$G/Z(G)$ is a Frobenius group with Frobenius kernel and complement $F/Z(G)$ and $K/Z(G)$, respectively and F and K are abelian subgroups of G; and $\omega(\mathcal{A}_G) = |F : Z(G)| + 1$.*
3. *$G/Z(G)$ is a Frobenius group with Frobenius kernel and complement $F/Z(G)$ and $K/Z(G)$, respectively; and K is an abelian subgroup of G, $Z(F) = Z(G)$, and $F/Z(G)$ is of prime power order; and $\omega(\mathcal{A}_G) = |F : Z(G)| + \omega(\mathcal{A}_F)$.*
4. *$G/Z(G) \cong S_4$ and V is a non-abelian subgroup of G such that $V/Z(G)$ is the Klein 4-group of $G/Z(G)$; and $\omega(\mathcal{A}_G) = 13$.*
5. *$G = A \times P$, where A is an abelian subgroup and P is an AC-subgroup of prime power order.*

Proof of Theorems 1.1 and 1.2. Let $q_1 = p_1^{n_1} > 3$ and $q_2 = p_2^{n_2} \geq 2$, where p_1 and p_2 are two prime numbers. Let $M_1 = \mathrm{GL}(2, q_1)$ and $M_2 = \mathrm{SL}(2, q_2)$ and suppose that G_1 and G_2 are two groups such that $\mathcal{A}_{G_i} \cong \mathcal{A}_{M_i}$ for $i = 1, 2$.
If $q_1 = 2$, then $M_2 \cong S_3$ is the symmetric group of degree 3 and so by Proposition 3.2 of [1], $G_2 \cong M_2$. If $q_2 = 3$, then M_2 is a group of order 24 and its center has order 2. As there is some element g with $|C_{G_2}(g)| = 6$, we see that there is no normal Sylow 3-subgroup in G_2. Hence $G_2/Z(G_2) \cong A_4$. So either $G_2 \cong M_2$ or $\mathbb{Z}_2 \times A_4$. But as there are elements $h \in G_2$ with $|C_{G_2}(h)| = 4$, we have $G_2 \cong M_2$.
Now let $q_2 > 3$. If q_2 is even, then $\mathrm{PSL}(2, q_2) \cong M_2$ and so $\mathcal{A}_{G_2} \cong \mathcal{A}_{\mathrm{PSL}(2,q_2)}$. Then by Corollary 5.3 of [1], $G_2 \cong \mathrm{PSL}(2, q_2) \cong M_2$. Therefore we may assume that $q_2 \geq 5$ is odd.
By Proposition 4.3 of [1], $|G_i| = |M_i|$ for $i = 1, 2$. By Lemma 3.5 of [1], M_i's are AC-groups and so by Lemma 2.1(2) G_i's are also AC-groups. Now since $\mathcal{A}_{G_i} \cong \mathcal{A}_{M_i}$ and $|G_i| = |M_i|$, by Lemma 2.1 we have the following equality between multisets

$$W_i = \{|C_{G_i}(x)| \mid x \in G_i \backslash Z(G_i)\} = \{|C_{M_i}(g)| \mid g \in M_i \backslash Z(M_i)\}, \ \ i = 1, 2.$$

Also, since the order of two graphs $\mathcal{A}_{G_i}$ and $\mathcal{A}_{M_i}$ are the same, we have that $|G_i| - |Z(G_i)| = |M_i| - |Z(M_i)|$ and so $|Z(G_i)| = |Z(M_i)|$ $(i = 1, 2)$. Therefore, it follows from Propositions 3.14 and 3.26 of [1] that the multiset W_1 (resp., W_2) consists of three distinct integers $(q_1 - 1)^2$, (resp., $(q_2 - 1)/2$) $q_1^2 - 1$ (resp., $(q_2 + 1)/2$) and $q_1(q_1 - 1)$ (resp., q_2) with multiplicities $\frac{q_i(q_i+1)}{2}$, $\frac{q_i(q_i-1)}{2}$ and $q_i + 1$, respectively.

We claim that both groups G_1 and G_2 are not nilpotent. Suppose, for a contradiction, that G_i is nilpotent, then so is $G_i/Z(G_i)$. Therefore $G_i/Z(G_i)$ has only one Sylow p_i-subgroup. Since W_1 (resp., W_2) contains $q_i + 1$ elements all equal to $q_1(q_1 - 1)$ (resp., q_2), there exist two non-central elements x_1 and y_1 in G_1 (resp., x_2 and y_2 in G_2) such that $C_{G_1}(x_1) \neq C_{G_1}(y_1)$ and $|C_{G_1}(x_1)| = |C_{G_1}(y_1)| = q_1(q_1 - 1)$ (resp., $C_{G_2}(x_2) \neq C_{G_2}(y_2)$ and $|C_{G_2}(x_2)| = |C_{G_2}(y_2)| = 2q_2$). Since $C_{G_i}(x_i)/Z(G_i)$ and $C_{G_i}(y_i)/Z(G_i)$ are of the same order q_i, they are Sylow p_i-subgroups of $G_i/Z(G_i)$. It follows that $C_{G_i}(x_i)/Z(G_i) = C_{G_i}(y_i)/Z(G_i)$ and so $C_{G_i}(x_i) = C_{G_i}(y_i)$, a contradiction.

Now we prove that both G_1 and G_2 cannot be solvable. Suppose, for a contradiction, that G_i's are solvable. Then since G_i are not nilpotent, it follows from Theorem 2.3 that G_i's satisfy one of properties (1)-(4) in Theorem 2.3. Since $q_i > 3$ is a prime power and q_2 is odd, both of $|G_1/Z(G_1)| = q_1(q_1^2-1)$ and $|G_2/Z(G_2)| = \frac{q_2(q_2^2-1)}{2}$ cannot equal to $|S_4| = 24$. Therefore G_i's do not satisfy (4). If G_i satisfies either (1) or (2), then W_i contains only two distinct elements, since in the case (1), if $x \in N\backslash Z(G_i)$, then $C_{G_i}(x) = N$; and if $x \in G\backslash N$ then $C_N(x) = Z(G_i)$; so $|C_{G_i}(x)| \in \big\{|G_i : N||Z(G_i)|, |N|\big\}$ for every non-central element $x \in G_i$, and in the case (3), $|C_{G_i}(x)| \in \big\{|K|, |F|\big\}$. This is not possible, since W_i has exactly three distinct elements.
Finally, suppose that G_i satisfies (3). Note that $C_{G_i}(x) = C_F(x)$ for every non-central element $x \in F$ and $C_{G_i}(x)$ is equal to the conjugate of K which contains the non-central element x. It follows that the three distinct elements of the multiset $W_i' = \{w/|Z(G)| \mid w \in W_i\}$ are $|K/Z(G_i)|, r^k, r^\ell$, where $|F/Z(G)| = r^m$ and r is a prime number. This is impossible, since no two of the numbers q_1, q_1+1 or q_1-1 (resp., q_2, $(q_2+1)/2$ or $(q_2-1)/2$) can simultaneously be powers of the same prime.

Hence G_i's are finite non-solvable AC-groups. By Theorem 2.2, G_i's satisfy one of the conditions (1)-(4) stated in Theorem 2.2. If G_i satisfies (3), then as A_6 has self-centralizing elements of order 4 and 5, G_i contains two elements x_i, y_i such that $|\frac{C_{G_i}(x_i)}{Z(G_i)}| = 4$ and $|\frac{C_{G_i}(y_i)}{Z(G_i)}| = 5$. This implies that $q_1 \in \{4,5\}$ and $q_2 = 9$. Therefore $|G_1/Z(G_1)| = 4\cdot(4^2-1)$ or $5\cdot(5^2-1)$, which is impossible, since $|G_1/Z(G_1)| = |\mathrm{PSL}(2,9)| = \frac{9\cdot(9^2-1)}{2}$. Since $M_2 = \mathrm{SL}(2,9)$, $|Z(M_2)| = 2$. But 3 divides $Z(G_2)$ by Theorem 2.2, a contradiction.
If G_i satisfies (4), then as $\mathrm{PGL}(2,9)$ contains self-centralizing elements of order 8 and 10, G_i contains two elements t_i and s_i such that $|\frac{C_{G_i}(t_i)}{Z(G_i)}| = 8$ and $|\frac{C_{G_i}(s_i)}{Z(G_i)}| = 10$. It follows that $\{8,10\} \subset \{q_2, \frac{q_2-1}{2}, \frac{q_2-1}{2}\}$, which is a contradiction as q_2 is a prime power; and for $i=1$, it follows that $q_1 = 9$. Hence $|Z(M_1)| = 8$. But 3 divides $|Z(G_1)|$, a contradiction. Thus G_i does not satisfy both (3) and (4).
Now suppose that G_i satisfies either (1) or (2). The group $\mathrm{PGL}(2,r^m)$ (resp. $\mathrm{PSL}(2,r^m)$) has a partition $\mathcal{P}$ consisting of r^m+1 Sylow r-subgroups, $\frac{(r^m+1)r^m}{2}$ cyclic subgroups of order r^m-1 (resp. $\frac{r^m-1}{\gcd(2,r^m-1)}$) and $\frac{(r^m-1)r^m}{2}$ cyclic subgroups of order r^m+1 (resp. $\frac{r^m+1}{\gcd(2,r^m-1)}$) (see pp. 185-187 and p.193 of [2]). Now [6, (5.3.3) in p.112] states that if $x \in G_i\backslash Z(G_i)$, then $C_{G_i}(x_i)/Z(G_i)$ belongs to $\mathcal{P}$. Suppose that $G_i/Z(G_i) \cong \mathrm{PGL}(2,r^m)$ (resp. $\mathrm{PSL}(2,r^m)$). Thus there exist elements $g_{i1}, g_{i2}, g_{i3} \in G_i\backslash Z(G_i)$ such that $|C_{G_i}(g_{i1})|/|Z(G_i)| = r^m$, $|C_{G_i}(g_{i2})|/|Z(G_i)| = r^m-1$ (resp. $\frac{r^m-1}{\gcd(2,r^m-1)}$), $|C_{G_i}(g_{i3})|/|Z(G_i)| = r^m+1$ (resp. $\frac{r^m+1}{\gcd(2,r^m-1)}$).
Therefore, if $G_i/Z(G_i) \cong \mathrm{PGL}(2,r^m)$ (resp. $\mathrm{PSL}(2,r^m)$), then $\{q_1-1, q_1, q_1+1\} = \{r^m-1, r^m, r^m+1\}$ (resp. $\{\frac{r^m-1}{\gcd(2,r^m-1)}, r^m, \frac{r^m+1}{\gcd(2,r^m-1)}\}$ and $\{\frac{q_2-1}{2}, \frac{q_2+1}{2}, q_2\} = \{r^m-1, r^m, r^m+1\}$ (resp. $\{\frac{r^m-1}{\gcd(2,r^m-1)}, r^m, \frac{r^m+1}{\gcd(2,r^m-1)}\}$).
It follows that, if $G_2/Z(G_2) \cong \mathrm{PGL}(2,r^m)$ then $q_2 = r^m+1$, $\frac{q_2+1}{2} = r^m$ and $\frac{q_2-1}{2} = r^m-1$. Since $q_2 \geq 5$, we have a contradiction as $3 \leq q_2 - \frac{q_2-1}{2} = r^m+1-r^m+1 = 2$. Hence $G_2/Z(G_2) \cong \mathrm{PSL}(2,r^m)$, $G_2' \cong \mathrm{SL}(2,r^m)$ and $r^m = q_2$. Now since $|G_2'| = |G_2| = |M_2|$, we have that $G_2 \cong M_2 = \mathrm{SL}(2,q_2)$. This completes

the proof of Theorem 1.2.
Now if $G_1/Z(G_1) \cong \mathrm{PGL}(2, r^m)$ (resp. $\mathrm{PSL}(2, r^m)$), it follows that $q_1 = r^m$ (resp. $q_1 = 2^m$). Since $\mathrm{PSL}(2, 2^m) \cong \mathrm{PGL}(2, 2^m)$, we have if G_1 satisfies either (1) or (2), then $G_1/Z(G_1) \cong \mathrm{PGL}(2, q_1)$ and $G_1' \cong \mathrm{SL}(2, q_1)$.
Therefore G_1 is a group satisfying the following conditions:

$$G_1/Z(G_1) \cong \mathrm{PGL}(2, q_1) \ \ (\bullet), \ \ G_1' \cong \mathrm{SL}(2, q_1) \quad \text{and} \ \ |Z(G_1)| = q_1 - 1.$$

If $q_1 = 2^m$ for some integer $m > 1$, then $\mathrm{SL}(2, q_1) \cong \mathrm{PGL}(2, q_1) \cong \mathrm{PSL}(2, q_1)$. Thus as $\mathrm{PSL}(2, q_1)$ is a non-abelian simple group, it follows from $(\bullet)$ that $G_1 = G_1'Z(G_1)$; and since G_1' is also non-abelian simple, $G_1' \cap Z(G_1) = 1$. Therefore $G_1 = G_1' \times Z(G_1)$. This completes the proof of Theorem 1.1. □

Acknowledgments. This work was done during author's sabbatical leave study in Summer 2007 at ICTP, Trieste, Italy. He is grateful to University of Isfahan for its financial support as well as ICTP for their warm hospitality. He was also supported by the Center of Excellence for Mathematics, University of Isfahan. The author is indebted to the referee for his/her careful reading, valuable comments and pointing out a serious error in the previous version of Theorem 1.1.

References

- [1] A. Abdollahi, S. Akbari and H.R. Maimani, *Non-commuting graph of a group*, J. Algbera **298** (2006), 468-492.
- [2] B. Huppert, *Endliche Gruppen I*, Springer-Verlag, Berlin, 1967.
- [3] A. R. Moghaddamfar, W. J. Shi, W. Zhou and A. R. Zokayi, *On the noncommuting graph associated with a finite group*, Siberian Math. J. **46** (2005), no. 2, 325-332.
- [4] B.H. Neumann, *A problem of Paul Erdös on groups*, J. Austral. Math. Soc. Ser. A **21** (1976), 467-472.
- [5] L. Pyber, *The number of pairwise noncommuting elements and the index of the centre in a finite group*, J. London Math. Soc. (2) **35** (1987), no. 2, 287-295.
- [6] R. Schmidt, *Zentralisatorverbände endlicher gruppen*, Rend. Sem. Mat. Univ. Padova **44** (1970), 97-131.
- [7] L. Wang and W. Shi, *A new characterization of $L_2(q)$ by its noncommuting graph*, Front. Math. China **2(1)** (2007), 143-148.
- [8] L. Wang and W. Shi, *A new characterization of A_{10} by its noncommuting graph*, Comm. Algebra Comm. Algebra **36** (2008), no. 2, 523-528.